

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/349605083>

Resolving diurnal dynamics of the chloroplastic glutathione redox state in *Arabidopsis* reveals its photosynthetically-derived oxidation

Article in *The Plant Cell* · February 2021

DOI: 10.1093/plcell/koab068

CITATION

1

READS

189

6 authors, including:



Andreas Meyer

University of Bonn

221 PUBLICATIONS 7,093 CITATIONS

[SEE PROFILE](#)



Einat Zelinger

Hebrew University of Jerusalem

42 PUBLICATIONS 349 CITATIONS

[SEE PROFILE](#)



Matanel Hipsch

Hebrew University of Jerusalem

4 PUBLICATIONS 4 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Project

Zvika Hayuka [View project](#)



Project

Yael Heifetz [View project](#)

RESEARCH ARTICLE

Resolving diurnal dynamics of the chloroplastic glutathione redox state in *Arabidopsis* reveals its photosynthetically-derived oxidation

Zechariah Haber^a, Nardy Lampl^a, Andreas J. Meyer^b, Einat Zelinger^c, Matanel Hipsch^a and Shilo Rosenwasser^{a,d}

^a The Robert H. Smith Institute of Plant Sciences and Genetics in Agriculture, The Hebrew University of Jerusalem, Rehovot 7610000, Israel

^b Institute of Crop Science and Resource Conservation (INRES), Rheinische Friedrich-Wilhelms Universität Bonn, Friedrich-Ebert-Allee 144, D-53113 Bonn, Germany

^c The Interdepartmental Equipment Unit, The Robert H. Smith Faculty of Agriculture, Food & Environment, The Hebrew University of Jerusalem, Rehovot 7610001, Israel

^d Corresponding author: shilo.rosenwaser@mail.huji.ac.il

Short title: Diurnal dynamics of chloroplastic GSH redox state

One sentence summary: Monitoring the daily *in vivo* dynamics of the chloroplastic GSH redox state in light-stressed wild-type plants vs. photoprotective mutants provides insight into photosynthesis-dependent production of oxidants.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Shilo Rosenwasser (shilo.rosenwaser@mail.huji.ac.il)

ABSTRACT

Plants are subjected to fluctuations in light intensity, and this causes unbalanced photosynthetic electron fluxes and overproduction of reactive oxygen species (ROS). Electrons needed for ROS detoxification are drawn, at least partially, from the cellular glutathione (GSH) pool via the ascorbate-glutathione cycle. Here, we explore the dynamics of the chloroplastic glutathione redox potential (chl-E_{GSH}) using high-temporal-resolution monitoring of *Arabidopsis* (*Arabidopsis thaliana*) lines expressing the reduction-oxidation sensitive green fluorescent protein 2 (roGFP2) in chloroplasts. This was carried out over several days, under dynamic environmental conditions and in correlation with PSII operating efficiency. Peaks in chl-E_{GSH} oxidation during dark-to-light and light-to-dark transitions were observed. Increasing light intensities triggered a binary oxidation response, with a threshold around the light saturating point, suggesting two regulated oxidative states of the chl-E_{GSH}. These patterns were not affected in *npq1* plants, which are impaired in nonphotochemical quenching. Oscillations between the two oxidation states were observed under fluctuating light in WT and *npq1* plants, but not in *pgr5* plants, suggesting a role for PSI photoinhibition in regulating the chl-E_{GSH} dynamics. Remarkably, *pgr5* plants showed an increase in chl-E_{GSH} oxidation during the nights following light stresses, linking daytime photoinhibition and nighttime GSH metabolism. This work provides a systematic view of the dynamics of the *in vivo* chloroplastic glutathione redox state during varying light conditions.

Introduction

As sessile organisms that grow under highly variable light intensities, plants must constantly sense, respond, and adapt to the instantaneous modulation of daytime photon fluxes. This regulation is critical when the absorption of light energy, and the resulting production of NADPH and ATP, exceed the capacity of downstream reactions. This may occur either during the dark-to-light transition, under high light, or under fluctuating light intensities (Tikhonov, 2015; Kono and Terashima, 2014). Under these circumstances, over reduction of the photosynthetic electron transport chain can produce deleterious reactive oxygen species (ROS) in chloroplasts.

Several photoprotective mechanisms have evolved in plants to achieve homeostasis and optimal photosynthetic performance under high light and fluctuating environmental conditions, thereby protecting the photosynthetic machinery from light stresses. Among them are energy dissipation via heat, known as non-photochemical quenching (NPQ; Demmig-Adams, 1990; Horton and Ruban, 1992), cyclic electron flow (CEF; Arnon et al., 1954; Tagawa et al., 1963) and the water-water cycle (WWC; Mehler, 1951). Plants also cope with ROS overproduction by generating low molecular weight antioxidants such as glutathione (GSH) and ascorbate, as well as antioxidative enzymes such as ascorbate peroxidase (APX) and peroxiredoxins (Mittler et al., 2004; Takahashi and Badger, 2011; Kono and Terashima, 2014).

The rapid NPQ component, known as qE, is induced by increasing acidification of the thylakoid lumen and involves the activity of violaxanthin de-epoxidase (VDE), which catalyzes the conversion of violaxanthin to zeaxanthin (Niyogi et al., 1998). Analysis of the VDE null mutant (*npq1*), demonstrated that this mechanism increases plant tolerance to variation in light intensity by protecting photosystem II (PSII) under fluctuating light conditions in the field (Külheim et al., 2002).

CEF and the WWC play a role in balancing the stromal ATP/NADPH ratio according to requirements of primary metabolism and the need for protecting photosystem I (PSI) from photoinhibition (Avenson et al., 2005; Shikanai, 2007; Takahashi et al., 2009; Miyake, 2010). CEF and the WWC thereby serve as electron sinks under excessive light conditions (Ort and Baker, 2002; Asada, 1999). In higher plants, two CEF pathways are known to exist: the PROTON GRADIENT REGULATION 5 (PGR5)-dependent pathway, which is believed to be the main pathway in C3 plants, and the NADH dehydrogenase-like (NDH)-complex-dependent

pathway (Shikanai, 2016). While PGR5 is a key player in protecting PSI functionality by regulating CEF (Munekage et al., 2002; Suorsa et al., 2012; Suorsa et al., 2013), the mechanism for its action is not yet clear. When grown under high light or fluctuating light, the *Arabidopsis thaliana* (Arabidopsis) PGR5 null mutant (*pgr5*) shows stunted growth, impaired NPQ, a lower electron transport rate (ETR) relative to the wild type (WT) (Munekage et al., 2002) and also shows severe damage to PSI (Munekage et al., 2004; Suorsa et al., 2012). Therefore, exposure of *pgr5* to increased light intensities has served as an experimental model for assessing the physiological effects of PSI photoinhibition (Tiwari et al., 2016).

During the WWC, electrons are donated from PSI to molecular oxygen in the Mehler reaction, yielding superoxide radicals (O_2^-), which are then dismutated to molecular oxygen (O_2) and H_2O_2 in a reaction catalyzed by superoxide dismutase (SOD; Mehler, 1951; Rizhsky et al., 2003). Subsequent reduction of H_2O_2 to water is mediated mainly by ascorbate peroxidases (APXs) and peroxiredoxins (PRX; Awad et al., 2015). APX relies on electrons from reduced glutathione (GSH), channeled to APX through the ascorbate-glutathione cycle (Foyer and Noctor, 2011). In addition, ascorbate can also be directly regenerated by ferredoxin or by NAD(P)H via MDA reductase activity (Asada, 1999). Thus, electrons for the detoxification of WWC-derived H_2O_2 are drawn, at least partially, from the cellular glutathione pool. Notably, the production of H_2O_2 through the WWC can play a regulatory role, allowing communication between the photosynthetic light-harvesting reactions and downstream metabolic processes by transmitting oxidative signals to redox-regulated proteins (Dangoor et al., 2012; Eliyahu et al., 2015).

In the past fifteen years, redox-sensitive green fluorescent proteins (roGFPs) have allowed for *in vivo* monitoring of the glutathione redox potential (E_{GSH}) at high spatiotemporal resolution (Meyer et al., 2007; Rosenwasser et al., 2010; Nietzel et al., 2019). *In vitro* characterization of roGFP showed that roGFP reduction and oxidation are mediated by glutaredoxins (GRXs), which catalyze the reversible electron flow between glutathione and target proteins (Meyer et al. 2007; Gutscher et al., 2008; Trnka et al., 2020). In addition, the redox status of roGFP was not significantly affected by thioredoxin (Trx) or other redox-active compounds such as NADPH and ascorbate (Meyer et al. 2007; Gutscher et al. 2008). The pronounced preference of roGFP for E_{GSH} was further corroborated by using GSH biosynthesis inhibitors and *Arabidopsis* mutants compromised in GSH biosynthesis and reduction of glutathione disulfide

(GSSG) (Meyer et al. 2007; Meyer and Dick, 2010; Marty et al., 2019). Importantly, recording of the roGFP fluorescence and its dynamics appears to reflect the transmission of redox signals, ultimately leading to modulation of the redox state of native target proteins of GRXs (Meyer, 2008; Rosenwasser et al., 2014).

Due to their high sensitivity, reversibility and insensitivity to pH alterations in the physiological range (Schwarzländer et al., 2008), roGFP-based redox sensors are powerful tools for investigating redox dynamics in subcellular compartments. Recently, roGFP-expressing plants and algae have been tested under various biotic and abiotic stresses and were shown to develop organelle-specific roGFP oxidation patterns (Bratt et al., 2016; van Creveld et al., 2015; Mizrachi et al., 2019; Volpert et al., 2018; Rosenwasser et al., 2010; Nietzel et al., 2019).

The balance between ROS production and its scavenging by the chloroplastic GSH pool was suggested to play a significant role in sensing and responding to light stress because a disruption of this balance by elevating either GSH or ROS levels results in oxidative stress (Foyer et al., 1997; Karpinski et al., 1997; Creissen et al., 1999; Mullineaux et al., 2018). However, the diurnal dynamics of the chloroplastic glutathione redox state in response to changing light conditions have not been systematically explored. Here, we examined diurnal fluctuations in chloroplastic E_{GSH} under different light conditions by developing an automated system for monitoring the roGFP oxidation state throughout the day and night. We discovered unique patterns of chloroplast-specific E_{GSH} under normal, high-light and fluctuating light conditions. Moreover, by examining these patterns in *Arabidopsis* lines that are mutated in key proteins involved in photoprotective mechanisms, we revealed their interaction with the chloroplast GSH redox state.

Results

The chloroplastic E_{GSH} redox state is directly affected by photosynthetic electron transport

In order to map the *in vivo* temporal alterations in the chloroplastic E_{GSH} (chl- E_{GSH}) under ‘steady state’ and light treatment conditions, we used a chloroplast-targeted redox-sensitive GFP 2 sensor (chl-roGFP2; Fig. 1). Chloroplast targeting was achieved by using either the transketolase signal peptide (chl-TKTP-roGFP2, Schwarzländer et al., 2008) or a 2-Cys peroxiredoxin A signal peptide (chl-PRXaTP-roGFP2, this study),

as both target proteins to the chloroplast stroma (König et al., 2002; Schwarzländer et al., 2008). Chloroplast targeting was verified by the overlap of the chl-roGFP2 fluorescence signal with the chlorophyll fluorescence signal (Fig. 1A and Supp. Fig. 1). To monitor the response of the entire *Arabidopsis* shoot rosette to imposed redox changes, three-week-old plantlets with ubiquitous expression of chl-roGFP2 were examined. Whole-plant *in vivo* fluorescence images showed that external application of H₂O₂ caused an increased emission after excitation at 405 nm fluorescence intensity and decreased when excited at 465 nm (Fig. 1B). Conversely, treatment of plants with DTT caused a less intense signal with 405 nm excitation and increased brightness with 465 nm excitation. (Fig. 1B). Ratiometric analysis of the raw images (405nm/465nm) resulted in informative false-color images showing the oxidation and reduction of the probe after application of H₂O₂ and DTT, respectively (Fig. 1B), as well as a positive correlation between the concentration of H₂O₂ applied and chl-roGFP2 oxidation (Supp. Fig. 2). These results validated the sensitivity of chl-roGFP2 to redox alterations and are consistent with probe performance, previously observed in plant and mammalian cells (Dooley et al., 2004; Jiang et al., 2006; Meyer et al., 2007; Schwarzländer et al., 2008).

To examine the dependence of the redox state of chl-roGFP2 on photosynthetic activity, we treated plants with various concentrations of H₂O₂ to induce oxidation, and we subsequently measured the recovery of chl-roGFP2 from oxidation under different light intensities. Regardless of the applied H₂O₂ concentration, the chl-roGFP2 fluorescence ratio immediately declined, resulting in distinct concentration-dependent peaks (Fig. 1C). Surprisingly, the decline was light-dependent and became faster and more efficient with increasing light intensities (Fig. 1C). Almost full reduction of chl-roGFP2 was only achieved when plants were illuminated with high light (1000 μ E m⁻² s⁻¹). In this case, ratios for plants treated with 500 mM H₂O₂ approached resting levels with low fluorescence ratios after about 200 min (Fig. 1C). While light-dependent chl-roGFP2 reduction was observed following H₂O₂ treatment, probe oxidation was observed in plants exposed to one hour of high light conditions (700 μ mol m⁻² s⁻¹), but not in plants pretreated with 150 μ M 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU), as determined by whole-plant imaging (Fig. 1D) and quantitative fluorometry (Fig. 1E). Compared to control plants, a higher level of oxidation was observed in DCMU-treated plants after 3h under high-light (HL). Taken together, these results suggest that the chl-roGFP2 redox state is directly affected by photosynthetic activity.

Distinct patterns of chloroplastic E_{GSH} during a diurnal cycle

The observed relationship between photosynthesis activity and chl-roGFP2 redox state motivated us to systematically monitor its oxidation patterns under normal growth light and under changing light conditions. To this end, we developed an automated system that allows for continuous measurement of the degree of oxidation (OxD) of chl-roGFP2 and chlorophyll-fluorescence-derived photosynthetic parameters from many plants under dynamic environmental conditions (see Methods). Examination of chl-roGFP2 oxidation under typical laboratory growth conditions, over the entire day (16 h light at $120 \mu\text{mol m}^{-2} \text{ s}^{-1}$ /8 h dark), showed a stable chl-roGFP2 state during the day of approximately $\text{OxD} = 25\%$. Taking into account a stromal pH of 8, this OxD would correspond to an $E_{GSH} = -347 \text{ mV}$. These results are in agreement with previous calculations of chloroplastic E_{GSH} under steady-state conditions (Schwarzländer et al., 2008; Rosenwasser et al., 2010). The average daytime OxD was then taken as a ‘steady-state’ reference point and pronounced deviations in OxD that occurred during the transition from dark to light and vice versa were expressed as ΔOxD . A peak of oxidation with an increase in OxD of about 10% ($\Delta\text{OxD} = \sim 10\%$) occurred during dark-light transition (Fig. 2A).

Surprisingly, a very similar peak of transient oxidation with the same amplitude occurred during the light-dark transition. In this case, however, OxD did not return to the steady-state level but rather became even more reduced as $\Delta\text{OxD} = -13\%$. This reduction was then followed by gradual oxidation, which after about 6 h stabilized just below the steady-state daytime reference value (Fig. 2A). To better interpret chloroplast redox-metabolism-alterations during the nighttime, we estimated stromal NADPH availability using *Arabidopsis* plants expressing the recently developed NADPH sensor (TKTP-iNAP4; Tao et al., 2017; Lim et al., 2020). The observed increase in the TKTP-iNAP4 ratio (Fig. 2B), obtained by dividing the emissions at 510 nm after sequential excitation at 400 nm and 485 nm suggests an increase in NADPH levels at the beginning of the night, implying that the corresponding reduction of E_{GSH} is governed by the higher reduction rate of the GSH pool by Glutathione Reductase (GR).

To further characterize the oxidation occurring during dark-to-light transitions, we collected data from 20 experiments, cumulatively involving approximately 2600 plants. This large data set allowed for a high-temporal resolution analysis of the chl-roGFP2 OxD values that surround the transitions points. Within the first minute of the light phase, we observed a rapid burst in chl-roGFP2 OxD levels that remained stable

for 10-15 min, and this was followed by chl-roGFP2 reduction towards steady state day time values determined after about 20 min (Fig. 2C). Contrastingly, chl-roGFP2 oxidation in the light-dark transition occurred approximately 5 min after lights were switched off. This oxidation of $\Delta\text{OxD} = \sim 10\%$ was followed by a gradual reduction, reaching values that were detected during the day after 20 min (Fig. 2D). These results demonstrate significant oxidation of stromal *E_{GSH}* during darkness-light transitions and imply the transmission of oxidative signals during the induction and termination phases of photosynthesis.

The chloroplastic *E_{GSH}* dynamically responds to high-light conditions.

Next, we focused on investigating the dynamics of chl-*E_{GSH}* under HL conditions. First, we assessed the induction of the WWC and the ascorbate-glutathione cycle under HL conditions by quantifying the level of expression of chloroplast-targeted genes involved in these pathways and several established marker genes for oxidative stress (Fig. 2E, The complete list of genes is described in Supplemental Data Set 1). Plants were exposed to three hours of normal growth conditions ($120 \mu\text{mol m}^{-2} \text{s}^{-1}$) at the beginning of the day, followed by a phase of different HL conditions at 850 or $1700 \mu\text{mol m}^{-2} \text{s}^{-1}$ for a 6-h period. After the HL phase, the light was dimmed again to the normal intensity for an additional 7-h period, followed by 8 h of darkness. Samples for RNA extraction were collected at different time points during the HL phase, after returning to normal growth conditions and during the subsequent dark phase. Control plants were exposed to normal growth conditions of $120 \mu\text{mol m}^{-2} \text{s}^{-1}$ for the entire 16-hour light period.

We observed induction in four genes encoding for chloroplastic superoxide dismutase (FSD1, FSD2, FSD3 and CSD2) and the gene for stromal ascorbate peroxidase (sAPX) during the HL phase. Moderate induction was also observed for monodehydroascorbate reductase (MDAR6), and dehydroascorbate reductase (DHAR3) and two peroxiredoxins (PRXQ and PrxIIE). No changes in the expression level of glutathione reductase 2 (GR2) were observed. We also observed the induction of two H_2O_2 specific markers FER1, FIN4 (op den Camp et al., 2003; Gadjev et al., 2006) as early as 1 hour into the HL phase in both examined HL irradiances and the induction of two general oxidative stress markers, TI1 and At2g21640, mainly in response to $1700 \mu\text{mol m}^{-2} \text{s}^{-1}$. These results point to the induction of the WWC and ascorbate-glutathione cycle under the examined HL conditions.

We thus investigated chl-roGFP2 OxD patterns under HL conditions by exposing plants to various HL intensities during a 24-h cycle. Plants were exposed to three hours of normal growth light conditions ($120 \mu\text{mol m}^{-2} \text{s}^{-1}$) at the beginning of the day, after which the light irradiance was raised to various intensities (220, 330, 450, 550, 650, 750, 850, 1200, 1500 or $1700 \mu\text{mol m}^{-2} \text{s}^{-1}$) for a 6-h period. After the HL phase, the light was dimmed to the normal growth light conditions for an additional 7-h period, followed by 8 hours of darkness. (Fig. 2F inset). In addition to chl-roGFP2 oxidation (Fig. 2F), the PSII operating efficiency (Φ_{PSII}) was continuously monitored based on chlorophyll fluorescence imaging (Fig. 2G).

We observed immediate chl-roGFP2 oxidation, followed by a gradual reduction, when plants were shifted from $120 \mu\text{mol m}^{-2} \text{s}^{-1}$ to any higher light intensity, indicating a rapid chl-Egsh response to unpredicted increases in the light input (Fig. 2F). No significant changes in the TKTP-iNAP4 ratio were detected upon shifting of plants from normal growth light conditions to HL of $850 \mu\text{mol m}^{-2} \text{s}^{-1}$, implying that higher ROS production rates dominated the increase in chl-roGFP2 oxidation (Supp. Fig. 3). Further reduction and stabilization of the chl-roGFP2 redox state were observed when plants were returned to normal growth light conditions after the HL period (Fig. 2F). Reduction at the onset of darkness, followed by gradual oxidation, as observed during normal growth conditions, was observed for all light treatments (Fig. 2F). Similar chl-roGFP2 diurnal patterns were observed when plants were exposed to similar light treatments under elevated CO_2 levels (850 ppm), despite slightly higher reduction rates during the HL phase (Supp. Fig. 4). These light-induced chl-roGFP2 oxidation patterns were not observed in any of the DCMU-treated plants, which exhibited gradual oxidation of the roGFP2 probe, further pointing to the direct regulation of roGFP2 dynamics by photosynthesis (Supp. Fig. 5).

To ensure that the observed HL responses were not affected by the availability of endogenous GRXs, which catalyze the reversible electron flow between GSH and roGFP, we examined oxidation patterns in plants expressing chloroplast-targeted Grx1-roGFP. In this probe, the human glutaredoxin-1 (Grx1) is fused to roGFP2; hence its equilibrium with the glutathione redox couple is not dependent on endogenous GRXs (Marty et al., 2009; Park et al., 2013). Similar diurnal oxidation patterns to those observed with unfused roGFP2 under HL were detected using the Grx1-roGFP (Supp. Fig. 6). Furthermore, the examination of the expression level of seven genes coding for chloroplast targeted GRXs under HL conditions showed that their expression levels

were either unchanged or slightly upregulated (Supp. Fig. 7). These results suggest that both probes are suitable for monitoring the chloroplastic E_{GSH} under dynamic light conditions.

Examination of the correlation between chl-roGFP2 oxidation and light irradiance revealed a binary response rather than a quantitative correlation. Irradiance values of 220-650 $\mu\text{mol m}^{-2} \text{ s}^{-1}$ triggered a similar initial chl-roGFP2 ΔOxD of approximately 10%, which is equivalent to an increase in E_{GSH} of approximately 6 mV. In contrast, irradiances of 750-1700 $\mu\text{mol m}^{-2} \text{ s}^{-1}$, the initial oxidation was approximately 20%, which is equivalent to an increase in E_{GSH} of approximately 13mV. In contrast, the decrease in Φ_{PSII} correlated with the light intensity (Fig. 2G). This binary response suggests a light-dependent threshold of chl-roGFP2 oxidation near 750 $\mu\text{E m}^{-2} \text{ s}^{-1}$, which matched the light saturation area, as indicated by light-response curves generated based on chlorophyll fluorescence measurements (Supp. Fig. 8). Overall, the data show the differential response of photosynthesis efficiency and redox metabolism to HL and suggest a regulatory role of chl- E_{GSH} during the shift from light-unsaturated to light-saturated photosynthesis.

Fluctuating light triggers oscillations in stromal E_{GSH}

As plants in the field are inevitably confronted with varying light intensities, we further examined chl-roGFP2 redox dynamics under conditions of fluctuating light (FL) (Fig. 3). To this end, after an initial illumination phase at 120 $\mu\text{mol m}^{-2} \text{ s}^{-1}$, plants were exposed, in the middle of the day, for 6 hours of FL alternating between 1700 $\mu\text{mol m}^{-2} \text{ s}^{-1}$ and 120 $\mu\text{mol m}^{-2} \text{ s}^{-1}$, in intervals of 1, 5 or 10 min (Fig. 3A, D and G). Under all tested frequencies, a gradual and partial loss of photosynthetic activity was observed, as indicated by decreasing Φ_{PSII} values, reaching a minimum at the end of the FL period (Supp. Fig. 9). Shifting plants back to normal light conditions resulted in partial Φ_{PSII} recovery, with values still lower than those measured at the beginning of the day (Supp. Fig. 9). Plants exposed to the 1 min intervals showed a pattern that resembled HL treatments, including initial chl-roGFP2 oxidation ($\Delta\text{OxD} = \sim 20\%$, Fig. 3B), followed by a gradual reduction when plants were moved back to normal growth light conditions. In contrast, oscillations in the chl-roGFP2 oxidation were observed in plants exposed to intervals of 5 min and 10 min (Fig. 3E and H).

To make sense of the original data, we simplified the visual analysis and extracted all values recorded during the 6-hour FL treatment and combined these data

into a single apparent FL cycle, *i.e.*, all ΔOxD values were sorted according to their relative time in the FL cycle (*e.g.* from the 10 min FL cycle experiment depicted in Fig. 3H, values for both 20 min and 40 min were considered the 20th minute each for their respective cycle and thus pooled together). This analysis resulted in a surprisingly clear picture showing an oscillation between higher and lower chl-roGFP2 oxidation states during the 5 min and 10 min FL cycles, while no oscillations could be observed for the 1-min FL cycle (Fig. 3C, F, I). For the 5-min and 10-min FL cycles, we observed a gradual increase in oxidation during the HL phase and a decrease during the subsequent normal light phase. Importantly, these patterns were conserved in all FL cycles, and were not altered despite the gradual decrease in Φ_{PSII} during the FL phase (Supp. Fig. 9). It should be noted that the fact that no oscillations were observed in plants exposed to 1-min intervals does not necessarily mean that chl- E_{GSH} is constant under these conditions because equilibration of redox potentials between roGFP2 and the stromal glutathione pools involves a GRX-catalyzed biochemical reaction and is thus not instantaneous. Taken together, these results point to frequency-dependent oscillations of the chl- E_{GSH} under FL.

Damage of photosystem I suppresses oscillations of E_{GSH} in fluctuating light

To examine the cross-talk between chl- E_{GSH} patterns and light photoprotection mechanisms, the chl-roGFP2 probe was expressed in *npq1* and *pgr5* plants and chl-roGFP2 OxD was measured in parallel with Φ_{PSII} under the same HL regimens (Fig. 4). Despite the clear decrease in Φ_{PSII} in *npq1*, as compared to WT plants under identical HL conditions (*e.g.*, reaching approximately 0.3 and 0.5 under 1700 $\mu\text{mol m}^{-2} \text{ s}^{-1}$, for *npq1* and WT plants, respectively; Fig. 4D, E), nearly identical chl-roGFP2 oxidation trends were observed in both lines (Fig. 4A, B). Severely impaired photosynthetic efficiency was observed in *pgr5* plants exposed to HL, reaching Φ_{PSII} values of approximately 0.15 in plants illuminated at 1700 $\mu\text{mol m}^{-2} \text{ s}^{-1}$ (Fig. 4F). As in WT, shifting of *pgr5* to HL resulted in fast oxidation, but chl-roGFP2 OxD values recovered much quicker than in WT (Fig. 4C and Fig. 4A, respectively). These results suggest that photoinhibition of PSII does not necessarily result in stronger chl- E_{GSH} oxidation and that PSI photoinhibition results in faster reduction rates of chl- E_{GSH} .

As PGR5 was found to be essential for photoprotection, specifically under FL conditions (Yamamoto and Shikanai, 2019; Yamori et al., 2016; Munekage et al., 2002; Suorsa et al., 2013), we examined chl-roGFP2 oxidation in WT, *npq1* and *pgr5* plants

under FL conditions with 1- and 10-min intervals, as explained earlier. As shown for HL, FL also resulted in an apparent decrease in Φ_{PSII} in *npq1* and *pgr5* compared to WT (Supp. Fig. 10). Recording of Δ_{OxD} over an entire diurnal cycle with a FL phase of 1-min intervals did not significantly impact the overall oxidation pattern compared to plants exposed to continuous HL (Supp. Fig. 11A). Accordingly, no significant difference in oxidation patterns was observed between the HL phase and the normal-light phases of each cycle in all three lines (Supp. Fig. 12). Under FL condition of 10-min intervals, the chl-roGFP2 redox state oscillated between relatively reduced and relatively oxidized values, according to the relative timing within the FL cycle in WT and *npq1* plants (Fig. 4G and H). Intriguingly, the fluctuation in Δ_{OxD} during the FL cycle could not be observed in *pgr5* plants (Fig. 4I). Taken together, these results demonstrate that the chl- E_{GSH} response to FL conditions is dependent on PGR5 activity and is impaired under PSI photodamage.

Gradual relaxation of chloroplastic E_{GSH} oxidation and Φ_{PSII} impairment in *pgr5* plants

Interestingly, a significant increase in chl-roGFP2 OxD was observed in *pgr5* plants during the night following high and FL conditions, denoting an increase in chl- E_{GSH} (Fig. 4C and Supp. Fig. 11A-B). As shown, oxidation levels gradually increased during the night, reaching their maximum level at the end of the night period. The observed shift in OxD of approximately 25% (compared to the end-of-day values) in the night following HL phase of $1700 \mu\text{mol m}^{-2} \text{ s}^{-1}$ denote an increase in chl- E_{GSH} of approximately 16 mV. To ensure that the observed light response in *pgr5* is not affected by the availability of endogenous GRXs. We examined the expression level of chloroplast targeted GRXs at the end of the night. No significant changes in the expression level of most chloroplast targeted GRXs were observed in *pgr5* plants exposed to HL compared to those exposed to normal light or WT (Supp. Fig. 13), suggesting no alterations in the availability of endogenous GRXs at this time point. At the end of the night, chl-roGFP2 OxD values returned to steady state OxD levels immediately when the light was turned on. In agreement with the chl- E_{GSH} oxidation, we detected an induction of several oxidative stress marker genes (F6'H1, DLO1 and TI1) at the end of the night exclusively in *pgr5* plants exposed to HL (Supp. Fig. 13).

To further elucidate this phenomenon, we subjected *pgr5* plants to 72-h experiments consisting of 24-h HL experiments (containing a 6-h HL period of 850 or

1700 $\mu\text{mol m}^{-2} \text{s}^{-1}$), followed by two days with normal light (Fig. 5A, B; data *gll*, which served as the background line for the *pgr5* mutation, are presented in Fig. 5C, D). During the first day of recovery under normal-light conditions, Φ_{PSII} values were still lower than in *gll* plants (0.5–0.56 in *pgr5* versus approximately 0.7 in *gll*), and night chl-roGFP2 OxD oxidation clearly occurred again, albeit to a lower degree. Complete relaxation of the night chl-roGFP2 OxD oxidation was observed during the third night after HL treatment, reaching chl-roGFP2 OxD and Φ_{PSII} values comparable to those observed under normal light conditions (Supp. Fig. 14). During the days, despite night oxidation, the chl-roGFP2 values were comparable on all the examined days. This night oxidation effect, which appeared hours after the HL stress, suggests a link between photoinhibition during the day and alterations in redox metabolism during the nights following the stress conditions.

Discussion

The light-dependent flow of electrons from water to ferredoxin, the photosynthetic linear electron flow, produces the reducing power required for diverse downstream metabolic events, including carbon, nitrogen and sulfur assimilation, as well as for antioxidant activity. On the other hand, reduction of molecular oxygen at the acceptor site of PSI is the primary source of superoxide anion radical (O_2^-) and H_2O_2 in photosynthesizing chloroplasts (Asada, 1999; Badger et al., 2000). Thus, solar energy is the source for reducing power and oxidizing agents, subsequently requiring tight regulation to avoid imbalances between them.

The reduction of H_2O_2 via the ascorbate-glutathione pathway is thought to be the major mechanism leading to GSSG accumulation in plants (Rahantaniaina et al., 2013). Although the direct role of the GSH pool in detoxifying photosynthetically produced ROS has not been directly proven, and the involvement of DHARs has been debated (Rahantaniaina et al., 2017; Polle, 2001), a recent study demonstrated the role of GSH in recycling ascorbate under high-light conditions either nonenzymatically or via DHAR activity (Terai et al., 2020). Accordingly, light-dependent alterations in chl-EGSH likely reflect the balance between photosynthesis-dependent H_2O_2 production and NADPH availability for GR activity.

By applying whole-plant fluorescence imaging, we resolved the probe's spatial fluorescence patterns and validated probe expression in the entire tissue (Fig. 1B). This

is of interest as partial lack of the fluorescence signals resulting from gene silencing is frequently reported for plant protein-based biosensors (Jones et al., 2014; Exposito-Rodriguez et al., 2017). As silencing of the genetically encoded probe increase over generations, working with earlier generations of the desired line is typically recommended (Schwarzländer et al., 2016). Specifically, we have found chl-roGFP2 expression to be stable across generations in WT and the mutant lines used in this study. Moreover, our measurements with chl-roGFP2 provided the same readout regardless of whether or not a catalyzing GRX was fused to the probe or not. This suggests that the amounts of GRXs capable of catalyzing the redox equilibration of roGFP2 with E_{GSH} are sufficient in chloroplasts for reliable measurements of chl- E_{GSH} even with free roGFP2. It is generally assumed that endogenous GRX-target proteins with redox-sensitive thiols will respond to changes in chl- E_{GSH} in a similar form as roGFP2 (Meyer, 2008). Thus, changes in chl- E_{GSH} monitored with roGFP2 might indicate a light-induced modulation in the activity of such stromal proteins.

By measuring chl-roGFP2 oxidation at high temporal resolution over several consecutive days and in response to diverse light conditions, we revealed the dynamic response of chl- E_{GSH} to changing light conditions. The results presented showed oxidation peaks during darkness-light and light-darkness transitions (Fig. 2C, D). During the former, the discrepancy between the light-dependent reactions, which are activated immediately upon light exposure, and the Calvin-Benson-Bassham cycle reactions, which involves a 15-min activation phase, resulted in a limited overflow of electrons from PSI to NADP⁺ (Wirtz et al., 1982; Tikhonov, 2015) and transient diversion of electrons to O₂ reduction. The subsequent reduction of the chl-roGFP2 oxidation state may result from the full induction of CO₂ assimilation activity and a decrease in O₂ reduction in parallel with H₂O₂ detoxification. These results align with the suggested induction of the WWC during the induction phase of photosynthesis (Radmer and Kok, 1976; Miyake, 2010; Volpert et al., 2018).

The light-dependent increase in oxidants, which triggers changes in chl- E_{GSH} , may transmit oxidative signals to redox-sensitive proteins during the photosynthesis induction phase (Buchanan and Balmer, 2005; Rahantaina et al., 2013). Indeed, the reception of oxidative signals by chloroplastic atypical thioredoxins (which regulate NPQ induction and starch biosynthesis) was demonstrated shortly after illumination (Dangoor et al., 2012; Eliyahu et al., 2015). Similarly, peaks in oxidant production during light-dark transitions can play a regulatory role in oxidizing redox-sensitive

proteins at night, allowing transmission of reducing signals when the light is turned on again. Oxidative signals transmitted from H₂O₂ to target proteins, mediated by 2-Cys PRX at the onset of the dark period, have been demonstrated (Ojeda et al., 2018; Vaseghi et al., 2018; Eliyahu et al., 2015). Similarly, oxidation of key proteins in photosynthesis, including the ATP synthase CF1- γ subunit, fructose 1,6-bisphosphatase (FBPase) and sedoheptulose 1,7-bisphosphatase (SBPase), has been observed after the light was switched off (Yoshida et al., 2014). Dynamic response of the E_{GSH} upon light-to-dark transition was also recently reported in the model moss *Physcomitrella patens* (Müller-Schüssle et al., 2020). However, the molecular features and biophysical mechanism underlying the induction of these oxidation peaks during light-dark transitions, as well as the possible involvement of the GSH/GRXs pathway in oxidizing target proteins during the photosynthesis induction and termination phases, are still to be resolved.

As opposed to a relatively reduced ‘steady-state’ under normal growth conditions, chl-roGFP2 underwent considerable oxidation under HL conditions followed by gradual reduction (Fig. 2F). Interestingly, light irradiances of 220-650 $\mu\text{mol m}^{-2} \text{ s}^{-1}$ induced a lower chl-roGFP oxidation than irradiances of 750-1700 $\mu\text{mol m}^{-2} \text{ s}^{-1}$, while within the two ranges, there did not seem to be a significant difference. This binary effect, together with the contrasting nonbinary behavior of the drop in Φ_{PSII} during the HL period, suggests a regulatory role for chl-E_{GSH}, rather than a safety valve for energy dissipation through the WWC. Furthermore, it also implies the existence of a threshold near the light saturation point, which separates between two distinct states of oxidant production and, consequently, the chl-E_{GSH}. We hypothesize that these two redox states signal distinct metabolic responses, in accordance with the severity of the HL conditions. The similarity in chl-E_{GSH} OxD patterns between WT and *npql* plants, despite the significantly lower PSII efficiency in the mutant line (Fig. 4), further supports this signaling role of chl-E_{GSH} dynamics, as nonphotochemical quenching is a major mechanism for dissipating excess photons in higher plants (Osmond et al., 2000). The combination of increased antioxidant activity, as demonstrated by the induction of antioxidant genes (Fig. 2E) as well as enhanced consumption of reducing equivalents, may explain the gradual reduction observed during the HL phase (Fig. 2F).

A frequency-dependent shift between two oxidation states was observed under FL conditions (Fig. 3), suggesting its role in photosynthesis regulation under natural

conditions, in which the solar flux varies due to the diurnal cycle, canopy structure and varying cloud coverage. Interestingly, while a decrease in photosynthesis efficiency was observed during FL (Supp. Fig. 9), the ability to shift between the two chl-roGFP2 oxidation states was not impaired throughout the FL period (Fig. 3). The fact that the observed fluctuating oxidation patterns were not induced in *pgr5* plants may point towards a connection between PSI photoinhibition and rates of ROS production under these conditions. This link is further displayed by the faster reduction rates observed in *pgr5* plants (Fig. 4C). It has recently been demonstrated that *pgr5* plants are highly sensitive to HL-induced damage of iron–sulfur (Fe–S) clusters of PSI. This damage provides an additional photoprotective mechanism by inducing a nonphotochemical photoprotective energy quenching state (Tiwari et al., 2016). Based on the data observed here, we suggest that this PSI quenching state results in impaired regulation of the chloroplast EGSH. This results are in line with the proposed role of PSI photoinhibition in preventing excessive ROS production (Lima-Melo et al., 2019a) and the increased antioxidant capacity and decreased production of superoxide and H₂O₂ in *pgr5* plants (Suorsa et al., 2012). Accordingly, alterations in ROS production and downstream redox signaling may affect the sensitivity of *pgr5* to FL conditions.

While focusing on light-dependent oxidant production, continuous diurnal chl-roGFP2 OxD measurement uncovered a unique phenotype in *pgr5* plants during the nights following the HL and FL conditions. Pronounced gradual oxidation was observed on the first night following the HL stress, reaching the highest observed OxD from all experiments by the end of the night (Fig. 4C). Furthermore, nighttime oxidation was observed again the next night and typical night redox patterns seen in control plants were only observed in *pgr5* plants on the third night after treatment (Fig. 5A). This effect, which only surfaced hours after exposure to light stress, was not directly caused by photoinhibition and may be related to the slow recovery time of PSI (Zhang et al., 2011; Sonoike, 2011). Accordingly, restoration of CO₂ assimilation in HL-treated *pgr5* plants was achieved during the third day in normal light conditions and was attributed to the recovery of PSI parameters, such as maximum oxidizable P700 and maximal reduction state of ferredoxin (Lima-Melo et al., 2019b).

It is possible that chl-EGSH oxidation, observed during the two nights after HL stress, results from starch starvation caused by PSI photoinhibition in *pgr5* plants (Lima-Melo et al., 2019b). It has been reported that starch accumulates in chloroplasts during the day and is metabolized during the night as an NADPH source via the pentose

phosphate pathway (Weise et al., 2004; Kirchsteiger et al., 2009). Therefore, the observed nighttime oxidation in *pgr5* plants following HL stress may be due to a deficiency in NADPH resulting from starch deficiency. Accordingly, the contrasting effect observed in WT, in which reduction of chl-*E_{GSH}* was observed at the beginning of the night, likely results from higher NADPH availability during the night, as observed using the TKTP-iNAP4 sensor (Fig. 2B). However, the demand for NADPH to maintain the highly reduced state of chl-*E_{GSH}* during the night assumes GSH oxidation at night.

Interestingly, earlier work demonstrated that in addition to their role in thiol-redox maintenance, GSH and GRXs participate in assembling Fe–S clusters and transferring them to acceptor proteins, linking GSH and iron metabolism (Kumar et al., 2011; Rouhier et al., 2010; Mühlenhoff et al., 2010; Rouhier et al., 2008). Accordingly, repair and biogenesis of PSI after HL-induced damage of Fe–S clusters may require the function of GSH and GRXs. It is possible that the observed oxidation of chl-*E_{GSH}* several nights after HL or FL stress in *pgr5* plants (Fig. 4C, Fig. 5 and Supp. Fig. 11), in which PSI centers are susceptible to photoinhibition, mirrors the diversion of reduced GSH for the slow process of biogenesis and repair of PSI Fe–S clusters. In view of that, the gradual reduction and oxidation of chl-*E_{GSH}* at night, as observed during normal growth conditions as well as during HL and FL conditions in WT plants (Fig. 2F and Fig. 3B, E and H), may reflect the natural repair daily cycle of PSI. Whether this repair machinery operates exclusively at nights, or that a shortage of photosynthetically produced reducing power in the dark allowed to uncover this GSH requirement is not yet clear. An increasing demand for GSH may occur under chilling stress conditions, in which PSI is the major site of photoinhibition (Terashima et al., 1994), resulting in slow repair of PSI Fe–S clusters that will be further aggravated when glutathione reductase activity is suppressed (Shu et al., 2011). Further work will be required to characterize the molecular mechanisms regulating nighttime chl-*E_{GSH}* oxidation and the possible connection to PSI recovery.

In conclusion, by continuously monitoring roGFP2 oxidation patterns over several days, we provide a comprehensive view of photosynthetically related alteration in the chloroplastic *E_{GSH}* under various light conditions and uncover an uncharacterized link between PSI photoinhibition and nighttime GSH metabolism. Monitoring of compartment-specific redox metabolism under varying environmental conditions will

further expand the fundamental understanding of the role of redox signaling in light acclimation of higher plants.

Materials and Methods

Plant material, growth conditions and experimental setup

Arabidopsis thaliana WT (ecotype Columbia-0), *npq1* (CS3771, At1g08550, obtained from ABRC), and *pgr5* (EMS mutant line, At2g05620, obtained from Prof. T. Shikanai; lines were used throughout this research. Columbia ecotype *glabrous 1* (*gl-1*, At3g27920, obtained from Prof. T. Shikanai [Kyoto University]), which served as the reference ecotype for the *pgr5* line, was also used. Plants were sown on soil, placed at 4°C for two days to ensure uniform germination, and then grown under 16-h/8-h light/dark cycles with a photosynthetic photon flux density (PPFD) of 120 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (21°C, 60-70% RH, ambient CO₂) for two-three weeks. For roGFP and PAM analysis, plants were transferred to 12-well cell culture multiwell plates in solid peat plugs and the plugs were covered (with holes for plants) with black plastic to prevent autofluorescence. For 3-(3,4-dichlorophenyl)-1,1-dimethylurea (DCMU) experiments, 2-3-week-old plants were sprayed with 150 μM DCMU (D2425-100G, Sigma), kept for 1 hour in the dark and then exposed to high light (HL) conditions of 700 $\mu\text{mol m}^{-2} \text{s}^{-1}$.

In all experiments, 2-3-week-old plants were incubated in 21°C, 60-70% RH and ambient CO₂. The following light conditions were applied: Normal growth experiments were carried out under 16 hours light (120 $\mu\text{mol m}^{-2} \text{s}^{-1}$)/8 hours dark per 24-hour cycle. For HL experiments, plants were exposed to 3 hours of normal light conditions (120 $\mu\text{mol m}^{-2} \text{s}^{-1}$), and then light irradiance was elevated during 6 hours to the following HL intensities: 220, 330, 450, 550, 650, 750, 850, 1200, 1500 or 1700 $\mu\text{mol m}^{-2} \text{s}^{-1}$. After the high-light phase, plants were grown again under normal light conditions for an additional 7 hours and finally were under an 8-hour darkness period. For fluctuating light (FL) experiments, plants were exposed to 3 hours of 120 $\mu\text{mol m}^{-2} \text{s}^{-1}$, followed by 6 hours of fluctuating light between 1700 (HL) and 120 (normal light) $\mu\text{mol m}^{-2} \text{s}^{-1}$ in frequencies of 1,5 or 10 minutes (*i.e.* 1,5 or 10 minutes of 1700 $\mu\text{mol m}^{-2} \text{s}^{-1}$ followed by 1, 5 or 10 minutes 120 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and vice versa *etc.*). Afterwards, plants were grown again for 7 hours under 120 $\mu\text{mol m}^{-2} \text{s}^{-1}$ and finally under an 8-hour darkness period.

Generation of chl-roGFP2 expressing lines

Chloroplast targeting was achieved by using either the transketolase signal peptide (Schwarzländer et al., 2008) or a 2-Cys peroxiredoxin A (PRXa; Q96291) signal peptide (this study), both of which will target proteins to the chloroplast stroma (König et al., 2002; Schwarzländer et al., 2008). For the generation of the chl-roGFP2 line, the full gene sequence was chemically synthesized. The first 74 amino acids from PRXa were used as a signal peptide. The chl-roGFP2 gene was cloned into the plant cloning vector pART7 using *Xho*I and *Hind*III restriction enzymes. The whole construct including the CaMV 35S promoter and ocs terminator was then cloned into the binary vector pART27 using the restriction enzyme *Not*I. The pART27 plasmid, which contains the chl-roGFP2 construct, was transformed into GV3101 *Agrobacterium tumefaciens*. Transformation of *A. thaliana* was performed by floral dip (Clough and Bent, 1998). Transformed lines were selected based on kanamycin resistance and the chl-roGFP2 fluorescence signal.

Confocal microscopy

Images were acquired with a Leica TCS SP8 confocal system (Leica Microsystems) and the LAS X Life Science Software, while using a HC PL APO $\times 40/1.10$ objective. All images were acquired at a resolution of 4096×4096 -pixels. Images were acquired with emission at 500-520 nm following excitation at 488 nm for chl-roGFP2 fluorescence and emission at 670 nm following excitation at 488 nm for chlorophyll fluorescence. Merged images were generated using Fiji (Image J) software (Schneider et al., 2012).

chl-roGFP2 fluorescence measurements and analysis

Whole-plant chl-roGFP2 fluorescence imaging was detected using an Advanced Molecular Imager HT (Spectral Ami-HT, Spectral Instruments Imaging, LLC., USA), and AMIview software for image acquisition. This analysis was used to screen for fluorescent plants and for qualitative analysis. For chl-roGFP2 fluorescence detection, excitation using $405 \text{ nm} \pm 10 \text{ nm}$ or $465 \text{ nm} \pm 10 \text{ nm}$ LED light sources and $510 \text{ nm} \pm 20 \text{ nm}$ emission filter were used. Highly oxidized chl-ro-GFP2 produces much greater fluorescence at 405 nm compared to 465 nm, whereas highly reduced chl-ro-GFP2 shows a stronger signal at 465 nm. Therefore the 405/465 ratio is a reliable indicator of

the degree of oxidation. For chlorophyll detection, a $405\text{ nm} \pm 10$ LED light source and a $670\text{ nm} \pm 20$ nm emission filter used were.

Quantitative chl-roGFP2 analysis was mainly carried out by the fluorometer method (Rosenwasser et al., 2010) using a Tecan Spark® multimode microplate reader. For all plate reader experiments, roGFP2 fluorescence was excited at $400\text{ nm} \pm 10$ nm and $485\text{ nm} \pm 10$ nm, respectively, and emission recorded at $520\text{ nm} \pm 5$ nm. Chlorophyll autofluorescence was excited at $400\text{ nm} \pm 10$ and recorded in the $670\text{ nm} \pm 20$ nm band. For extended times, plants were grown in 12-well plates and placed in a growth chamber (Fytoscope FS-SI-4600, Photon Systems Instruments, Czech Republic) for illumination at defined light intensities. For recording of chl-roGFP2 fluorescence, the well plates were automatically inserted into the plate reader using a robot (KiNEDx KX.01467, Peak Analysis and Automation (PAA), Hampshire, UK) controlled by a self-compiled program (Overlord™, PAA).

For the plate-reader analysis, a 9-by-9-pixel matrix was defined for each well and chlorophyll autofluorescence was detected, to generate a chlorophyll mask. This mask was then used to select pixels, which returned a positive chlorophyll fluorescence signal. Only those pixels were subsequently considered for the roGFP analysis. For background correction, the average signal of WT plants without chl-roGFP2 was determined and the values for this background autofluorescence subtracted from the values detected in the chl-roGFP2 fluorescence analysis. A similar experimental setup was used to measure the fluorescence emitted from the TKTP-iNAP4 and TKTP-iNAPc expressing lines. The pH-corrected ratio for TKTP-iNAP4 was calculated by dividing the fluorescence ratio values obtained from the TKTP-iNAP4 line by those obtained from the TKTP-iNAPc line, according to Lim et al. (2020). The degree of oxidation (the relative quantity of oxidized roGFP proteins, OxD) of roGFP2 was calculated based on the fluorescence signal according to Equation 1 (Meyer et al., 2007):

Equation 1:

$$OxD_{roGFP2} = \frac{R - R_{\text{red}}}{(I_{485\text{min}}/I_{485\text{max}})(R_{\text{ox}} - R) + (R - R_{\text{red}})}$$

where R represents the ratio (400/485) at each point in the experiment, R_{red} represents the ratio under fully reduced conditions, R_{ox} represents the ratio under fully oxidized

conditions, $I_{485\text{min}}$ represents the fluorescence emitted at 520 nm when excited at 485 nm under fully oxidized conditions and $I_{485\text{max}}$ represents the fluorescence emitted at 520 nm when excited at 485 nm under fully reduced conditions. In order to be able to calculate chl-roGFP2 OxD, we treated chl-roGFP2 *Arabidopsis* plants and measured the 400/485 nm fluorescence ratio under fully oxidized and fully reduced conditions by using 1-4 M H_2O_2 and 100 mM DTT, respectively, by immersing plants in the respective solutions. A custom-written Matlab script (available on request) was used to allow a robust analysis of multiple files, which contained the measurements from each experiment. The scripts collect the roGFP values from each plant into one matrix. In addition, experimental metadata such as the start and endpoint, plant lines, light irradiance, temperature, relative humidity (RH, %) and CO_2 concentration were also collected and added to the output file. E_{GSH} values were calculated according to Schwarzlander et al. (2008).

Chlorophyll fluorescence measurements

The Φ_{PSII} of each plant was detected in 60-min intervals using a Walz PAM IMAGING PAM M-series IMAG-K7 (MAXI) Fluorometer. The electron transport rate (ETR) was calculated as the multiplication of Φ_{PSII} by the quantity of photons emitted by the light source (PAR), by the average percentage of the photons absorbed by the photosynthetic machinery (0.84), by the amount of photons allocated to PSII (0.5; assuming that PSI and PSII contain the same amount of reaction centers, therefore each photosystem receives 50% of the photon influx). Accordingly, $\text{ETR} = Y(\text{II}) * \text{PAR} * 0.84 * 0.5$. Full ETR light induction curves' data are in Supp. Fig. 8. A custom-written Matlab script was used to allow a robust analysis of multiple files (available upon request).

Gene expression analysis

RNA was extracted from 3-week old rosettes using the Spectrum Plant RNA Kit (STRN50; Sigma). For each time point, three (for the one and three hr in HL) or two (for later time points) independent biological replicates were analyzed. The extracted RNA was digested and cleaned with the TURBO DNase kit (Ambion) according to the manufacturer's instructions and was reverse transcribed to cDNA with the ThermoScript RT-PCR system (Invitrogen). The qPrimerDB (Lu et al., 2018) was used for selecting gene-specific primers for reverse-transcription quantitative-PCR (RT-

qPCR). The BioMark HD system was used for high-throughput qPCR, using standard fast cycling conditions and melt-curve analysis, following the manufacturer's instructions (Fluidigm). Cycle threshold (C_t) values for each reaction (which correspond to the logarithm of the fold change in gene expression) were used for analysis and expression data was calculated using the ddCT approach (Livak and Schmittgen, 2001). The ACT2 gene (At3g18780) was used as a reference for equalizing the levels of RNA.

Statistics

Values in graphs mainly represent means of 8 plants for chl-roGFP2 OxD and 12 plants for Φ PSII, and error bars represent the respective standard errors. For many experiments, 3-4 plates were analyzed, and for a clear presentation, a “sliding window” approach was taken, in which each data point represents the average of three-four plates (24-32 plants). Statistical significance was tested using a two-tail Student's t-test and analysis of variance (ANOVA) at 95% confidence level and was indicated by asterisks when shown (Supplemental File 1).

Accession numbers

Germplasm used included *npq1* (CS3771, At1g08550, obtained from ABRC), *pgr5* (EMS mutant line, At2g05620, obtained from Prof. T. Shikanai) and *gl1* (At3g27920, obtained from Prof. T. Shikanai), which served as the reference ecotype for the *pgr5* line. Accession numbers for genes used in gene expression analyses are available in Supplemental Data Set 1.

Supplemental Data

Supplemental Figure 1: Confocal images of the chl-PRX_a_{TP}-roGFP2 line.

Supplemental Figure 2: Dose-dependent chl-roGFP2 oxidation in response to H₂O₂.

Supplemental Figure 3: Changes in chl-roGFP2 oxidation and NADPH levels under high-light conditions.

Supplemental Figure 4: Diurnal changes in chl-roGFP2 OxD and Φ PSII under high light and elevated CO₂.

Supplemental Figure 5: Changes in chl-roGFP2 oxidation in DCMU-treated plants under high-light conditions.

Supplemental Figure 6: Diurnal changes in chloroplast-targeted Grx1-roGFP2 oxidation under high-light conditions.

Supplemental Figure 7: Temporal changes in expression levels of chloroplast-targeted GRXs under high-light conditions.

Supplemental Figure 8: ETR light-response curves in WT, *npq1* and *pgr5* plants.

Supplemental Figure 9: Diurnal changes in Φ PSII during illumination with fluctuating light.

Supplemental Figure 10: Diurnal changes in Φ PSII during fluctuating light experiments in WT, *npq1* and *pgr5* plants.

Supplemental Figure 11: Diurnal changes in chl-roGFP2 OxD during fluctuating light experiments in WT, *npq1* and *pgr5* plants.

Supplemental Figure 12: Changes in chl-roGFP2 OxD during fluctuating light cycles in WT, *npq1* and *pgr5* plants.

Supplemental Figure 13: Temporal changes in expression levels of chloroplast-targeted GRXs, antioxidant genes and ROS gene markers in WT and *pgr5* plants at the end of night.

Supplemental Figure 14: Diurnal changes in chl-roGFP2 OxD and Φ PSII during the third day of recovery-from-high-light experiments in *pgr5* plants.

Supplemental Figure 15: Diurnal changes in chl-roGFP2 OxD during high-light experiments in wild type (WT) plants.

Supplemental Figure 16: Diurnal changes in chl-roGFP2 OxD during high-light experiments in wild type (WT) plants.

Supplemental Figure 17: Diurnal changes in chl-roGFP2 fluorescence ratios under high-light conditions in WT and mutants with defects in photoprotective mechanisms.

Supplemental Data Set 1. Temporal changes in the expression levels of chloroplast-targeted genes encoding antioxidant proteins, GRXs and established ROS markers in response to high-light conditions.

Supplemental File 1. Statistical analysis tables.

Author contributions: Z.H. and S.R. designed research. Z.H., N.L., E.L., M.H., and S.R. performed research. Z.H., and S.R. analyzed data. A.J.M. contributed new plant lines. Z.H., and S.R. wrote the manuscript with inputs from N.L. and A.J.M.

Acknowledgments: We thank Avihai Danon, Zach Adam and Assaf Vardi for critical comments on the manuscript. We thank Prof. Toshiharu Shikanai, who kindly provided the *pgr5* and *gl1* seeds for this research. We thank Prof. Wallace Boon Leong Lim, who kindly provided the TKTP-iNAP4 and TKTP-iNAPc *Arabidopsis* lines and Ms. Charis Chan for her help in shipping the seeds. We thank Daniel Waiger for his help in

acquiring the confocal images. We thank David Pilzer for his support in performing the gene expression analysis. This research was supported by the Israel Science Foundation (grant No. 826/17 and No. 827/17) to SR.

References

Arnon DI, Allen MB, Whatley FR (1954) Photosynthesis by isolated chloroplasts. *Nature* **174**: 394-396

Asada K (1999) The water-water cycle in chloroplasts: scavenging of active oxygens and dissipation of excess photons. *Annu Rev Plant Physiol Plant Mol Biol* **50**: 601–639

Avenson TJ, Cruz JA, Kanazawa A, Kramer DM (2005) Regulating the proton budget of higher plant photosynthesis. *Proc Natl Acad Sci U S A* **102**: 9709–9713

Awad J, Stotz HU, Fekete A, Krischke M, Engert C, Havaux M, Berger S, Mueller MJ (2015) 2-Cysteine peroxiredoxins and thylakoid ascorbate peroxidase create a water-water cycle that is essential to protect the photosynthetic apparatus under high light stress conditions. *Plant Physiol* **167**: 1592–1603

Badger MR, von Caemmerer S, Ruuska S, Nakano H (2000) Electron flow to oxygen in higher plants and algae: rates and control of direct photoreduction (Mehler reaction) and rubisco oxygenase. *Philos Trans R Soc Lond B Biol Sci* **355**: 1433-1446

Bratt A, Rosenwasser S, Meyer A, Fluhr R (2016) Organelle redox autonomy during environmental stress. *Plant Cell Environ* **39**: 1909–1919

Buchanan BB, Balmer Y (2005) Redox regulation: a broadening horizon. *Annu Rev Plant Biol* **56**: 187–220

Clough SJ, Bent AF (1998) Floral dip: a simplified method for *Agrobacterium* -mediated transformation of *Arabidopsis thaliana*. *Plant J* **16**: 735–743

Creissen G, Firmin J, Fryer M, Kular B, Leyland N, Reynolds H, Pastori G, Wellburn F, Baker N, Wellburn A, Mullineaux P (1999) Elevated glutathione biosynthetic capacity in the chloroplasts of transgenic tobacco plants paradoxically causes increased oxidative stress. *Plant Cell* **11**: 1277-1291

van Creveld SG, Rosenwasser S, Schatz D, Koren I, Vardi A (2015) Early perturbation in mitochondria redox homeostasis in response to environmental stress predicts cell fate in diatoms. *ISME J* **9**: 385–395

Dangoor I, Peled-Zehavi H, Wittenberg G, Danon A (2012) A chloroplast light-regulated oxidative sensor for moderate light intensity in *Arabidopsis*. *Plant Cell* **24**: 1894–1906

Demmig-Adams B (1990) Carotenoids and photoprotection in plants a role for the xanthophyll zeaxanthin. *Biochim Biophys Acta* **1020**: 1-24

Dooley CT, Dore TM, Hanson GT, Jackson WC, Remington SJ, Tsien RY (2004) Imaging dynamic redox changes in mammalian cells with green fluorescent protein indicators. *J Biol Chem* **279**: 22284–22293

Eliyahu E, Rog I, Inbal D, Danon A (2015) ACHT4-driven oxidation of APS1 attenuates starch synthesis under low light intensity in *Arabidopsis* plants. *Proc Natl Acad Sci* **112**: 12876–12881

Exposito-Rodriguez M, Laissue PP, Yvon-Durocher G, Smirnoff N, Mullineaux PM (2017) Photosynthesis-dependent H₂O₂ transfer from chloroplasts to nuclei provides a high-light signalling mechanism. *Nat Commun* **8**: 49

Foyer CH, Lopez-Delgado H, Dat JF, Scott IM (1997) Hydrogen peroxide- and glutathione-associated mechanisms of acclimatory stress tolerance and signalling. *Physiol Plant* **100**: 241–257

Foyer CH, Noctor G (2011) Ascorbate and glutathione: the heart of the redox hub. *Plant Physiol* **155**: 2–18

Gadjev I, Vanderauwera S, Gechev TS, Laloi C, Minkov IN, Shulaev V, Apel K, Inzé D, Mittler R, Breusegem FV (2006) Transcriptomic footprints disclose specificity of reactive oxygen species signaling in *Arabidopsis*. *Plant Physiol* **141**: 436–445

Gutscher M, Pauleau A-L, Marty L, Brach T, Wabnitz GH, Samstag Y, Meyer AJ, Dick TP (2008) Real-time imaging of the intracellular glutathione redox potential. *Nat Methods* **5**: 553–559

Hanson GT, Aggeler R, Oglesbee D, Cannon M, Capaldi RA, Tsien RY, Remington SJ (2004) Investigating mitochondrial redox potential with redox-sensitive green fluorescent protein indicators. *J Biol Chem* **279**: 13044–13053

Horton P, Ruban AV (1992) Regulation of photosystem II. *Photosynth Res* **34**: 375–385

Jiang K, Schwarzer C, Lally E, Zhang S, Ruzin S, Machen T, Remington SJ, Feldman L (2006) Expression and characterization of a redox-sensing green fluorescent protein (reduction-oxidation-sensitive green fluorescent protein) in *Arabidopsis*. *Plant Physiol* **141**: 397–403

Jones AM, Danielson JÅH, ManojKumar SN, Lanquar V, Grossmann G, Frommer WB (2014) Abscisic acid dynamics in roots detected with genetically encoded FRET sensors. *eLife* **3**: e01741

Karpinski S, Escobar C, Karpinska B, Creissen G, Mullineaux PM (1997) Photosynthetic electron transport regulates the expression of cytosolic ascorbate peroxidase genes in *Arabidopsis* during excess light stress. *Plant Cell* **9**: 627–640

Kirchsteiger K, Pulido P, González M, Cejudo FJ (2009) NADPH thioredoxin reductase c controls the redox status of chloroplast 2-cys peroxiredoxins in *Arabidopsis thaliana*. *Mol Plant* **2**: 298–307

König J, Baier M, Horling F, Kahmann U, Harris G, Schürmann P, Dietz K-J (2002) The plant-specific function of 2-Cys peroxiredoxin-mediated detoxification of peroxides in the redox-hierarchy of photosynthetic electron flux. *Proc Natl Acad Sci* **99**: 5738–5743

Kono M, Terashima I (2014) Long-term and short-term responses of the photosynthetic electron transport to fluctuating light. *J Photochem Photobiol B Biol* **137**: 89–99

Külheim C, Ågren J, Jansson S (2002) Rapid regulation of light harvesting and plant fitness in the field. *Science* **297**: 91–93

Kumar C, Igbaria A, D'Autreux B, Planson A-G, Junot C, Godat E, Bachhawat AK, Delaunay-Moisan A, Toledano MB (2011) Glutathione revisited: a vital function in iron metabolism and ancillary role in thiol-redox control. *EMBO J* **30**: 2044–2056

Lim S-L, Voon CP, Guan X, Yang Y, Gardeström P, Lim BL (2020) *In planta* study of photosynthesis and photorespiration using NADPH and NADH/NAD⁺ fluorescent protein sensors. *Nat Commun* **11**: 3238

Lima-Melo Y, Alencar VTCB, Lobo AKM, Sousa RH V, Tikkkanen M, Aro E-M, Silveira JAG, Gollan PJ (2019a) Photoinhibition of photosystem I provides oxidative protection during imbalanced photosynthetic electron transport in *Arabidopsis thaliana*. *Front Plant Sci* **10**: 916

Lima-Melo Y, Gollan PJ, Tikkkanen M, Silveira JAG, Aro E-M (2019b) Consequences of photosystem-I damage and repair on photosynthesis and carbon use in *Arabidopsis thaliana*. *Plant J* **97**: 1061–1072

Livak KJ. and Schmittgen, TD. (2001). Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta CT}$ Method. *Methods* **25**: 402–408.

Lu K, Li T, He J, Chang W, Zhang R, Liu M, Mengna Y, Fan Y, Ma J, Sun W et al (2018) qPrimerDB: a thermodynamics-based gene-specific qPCR primer database for 147 organisms. *Nucleic Acids Res* **46**: D1229-D1236

Marty L, Bausewein D, Müller C, Bangash SAK, Moseler A, Schwarzländer M, Müller-Schüssele SJ, Zechmann B, Riondet C, Balk J (2019) *Arabidopsis* glutathione reductase 2 is indispensable in plastids, while mitochondrial glutathione is safeguarded by additional reduction and transport systems. *New Phytol* **224**: 1569–1584.

Marty L, Siala W, Schwarzländer M, Fricker MD, Wirtz M, Sweetlove LJ, Meyer Y, Meyer AJ, Reichheld J-P, Hell R (2009) The NADPH-dependent thioredoxin system constitutes a functional backup for cytosolic glutathione reductase in *Arabidopsis*. *Proc Natl Acad Sci U S A* **106**: 9109–9114

Mehler AH (1951) Studies on reactions of illuminated chloroplasts: I. Mechanism of the reduction of oxygen and other hill reagents. *Arch Biochem Biophys* **33**: 65–77

Meyer AJ (2008) The integration of glutathione homeostasis and redox signaling. *J Plant Physiol* **165**: 1390–1403

Meyer AJ, Brach T, Marty L, Kreye S, Rouhier N, Jacquot J-P, Hell R (2007) Redox-sensitive GFP in *Arabidopsis thaliana* is a quantitative biosensor for the redox potential of the cellular glutathione redox buffer. *Plant J* **52**: 973–986

Meyer AJ, Dick TP (2010) Fluorescent protein-based redox probes. *Antioxid Redox Signal* **13**: 621–650

Mittler R, Vanderauwera S, Gollery M, Van Breusegem F (2004) Reactive oxygen gene network of plants. *Trends Plant Sci* **9**: 490–498

Miyake C (2010) Alternative electron flows (water–water cycle and cyclic electron flow around PSI) in photosynthesis: molecular mechanisms and physiological functions. *Plant Cell Physiol* **51**: 1951–1963

Mizrahi A, Graff van Creveld S, Shapiro OH, Rosenwasser S, Vardi A (2019) Light-dependent single-cell heterogeneity in the chloroplast redox state regulates cell fate in a marine diatom. *eLife* **8**: e47732

Mühlenhoff U, Molik S, Godoy JR, Uzarska MA, Richter N, Seubert A, Zhang Y, Stubbe J, Pierrel F, Herrero E, et al (2010) Cytosolic monothiol glutaredoxins function in intracellular iron sensing and trafficking via their bound iron-sulfur cluster. *Cell Metab* **12**: 373–385

Müller-Schüssle SJ, Wang R, Gütle DD, Romer J, Rodriguez-Franco M, Scholz M, Buchert F, Lüth VM, Kopriva S, Dörmann P (2020) Chloroplasts require glutathione reductase to balance reactive oxygen species and maintain efficient photosynthesis. *Plant J* **103**: 1140–1154

Mullineaux PM, Exposito-Rodriguez M, Laissac PP, Smirnoff N (2018) ROS-dependent signalling pathways in plants and algae exposed to high light: comparisons with other eukaryotes. *Free Radic Biol Med* **122**: 52–64

Munekage Y, Hashimoto M, Miyake C, Tomizawa K-I, Endo T, Tasaka M, Shikanai T (2004) Cyclic electron flow around photosystem I is essential for photosynthesis. *Nature* **429**: 579–582.

Munekage Y, Hojo M, Meurer J, Endo T, Tasaka M, Shikanai T (2002) PGR5 is involved in cyclic electron flow around photosystem I and is essential for photoprotection in *Arabidopsis*. *Cell* **110**: 361–371

Nietzel T, Elsässer M, Ruberti C, Steinbeck J, Ugalde JM, Fuchs P, Wagner S, Ostermann L, Moseler A, Lemke P, et al (2019) The fluorescent protein sensor roGFP2-Orp1 monitors *in vivo* H₂O₂ and thiol redox integration and elucidates intracellular H₂O₂ dynamics during elicitor-induced oxidative burst in *Arabidopsis*. *New Phytol* **221**: 1649–1664

Niyogi KK, Grossman AR, Björkman O (1998) Arabidopsis mutants define a central role for the xanthophyll cycle in the regulation of photosynthetic energy conversion. *Plant Cell* **10**: 1121– 1134

Ojeda V, Pérez-Ruiz JM, Cejudo FJ (2018) 2-Cys peroxiredoxins participate in the oxidation of chloroplast enzymes in the dark. *Mol Plant* **11**: 1377–1388

op den Camp RGL, Przbyla D, Ochsenbein C, Laloi C, Kim C, Danon A, Wagner D, Hideg É, Göbel C, Feussner I et al (2003) Rapid induction of distinct stress responses after the release of singlet oxygen in *Arabidopsis*. *Plant Cell* **15**: 2320-2332

Ort DR, Baker NR (2002) A photoprotective role for O₂ as an alternative electron sink in photosynthesis. *Curr Opin Plant Biol* **5**: 193–198

Park S-W, Li W, Viehhauser A, He B, Kim S, Nilsson AK, Andersson MX, Kittle JD, Ambavaram MMR, Luan S, et al (2013) Cyclophilin 20-3 relays a 12-oxo-phytodienoic acid signal during stress responsive regulation of cellular redox homeostasis. *Proc Natl Acad Sci U S A* **110**: 9559-9564

Polle A (2001) Dissecting the superoxide dismutase-ascorbate-glutathione-pathway in chloroplasts by metabolic modeling. Computer simulations as a step towards flux analysis. *Plant Physiol* **126**: 445-462

Radmer RJ, Kok B (1976) Photoreduction of O₂ primes and replaces CO₂ assimilation. *Plant Physiol* **58**: 336–340

Rahantaina M-S, Li S, Chatel-Innocenti G, Tuzet A, Mhamdi A, Vanacher H, Noctor G (2017) Glutathione oxidation in response to intracellular H₂O₂: key but overlapping roles for dehydroascorbate reductases. *Plant Signal Behav* **12**: e1356531

Rahantaina M-S, Tuzet A, Mhamdi A, Noctor G (2013) Missing links in understanding redox signaling via thiol / disulfide modulation : how is glutathione oxidized in plants. *Front Plant Sci* **4**: 1–13

Rizhsky L, Liang H, Mittler R (2003) The water-water cycle is essential for chloroplast protection in the absence of stress. *J Biol Chem* **278**: 38921–38925

Rosenwasser S, Rot I, Meyer AJ, Feldman L, Jiang K, Friedman H (2010) A fluorometer-based method for monitoring oxidation of redox-sensitive GFP (roGFP) during development and extended dark stress. *Physiol Plant* **138**: 493–502

Rouhier N, Couturier J, Johnson MK, Jacquot J-P (2010) Glutaredoxins: roles in iron homeostasis. *Trends Biochem Sci* **35**: 43–52

Rouhier N, Lemaire SD, Jacquot J-P (2008) The role of glutathione in photosynthetic organisms: emerging functions for glutaredoxins and glutathionylation. *Annu Rev Plant Biol* **59**: 143–166

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH image to ImageJ: 25 years of image analysis. *Nat Methods* **9**: 671-675

Schwarzländer M, Fricker MD, Müller C, Marty L, Brach T, Novak J, Sweetlove LJ, Hell R, Meyer AJ (2008) Confocal imaging of glutathione redox potential in living plant cells. *J Microsc* **231**: 299–316

Shikanai T (2007) Cyclic electron transport around photosystem I: genetic approaches. *Annu Rev Plant Biol* **58**: 199–217

Shikanai T (2016) Regulatory network of proton motive force: contribution of cyclic electron transport around photosystem I. *Photosynth Res* **129**: 253–260

Shu D-F, Wang L-Y, Duan M, Deng Y-S, Meng Q-W (2011) Antisense-mediated depletion of tomato chloroplast glutathione reductase enhances susceptibility to chilling stress. *Plant Physiol Biochem* **49**: 1228–1237

Sonoike K (2011) Photoinhibition of photosystem I. *Physiol Plant* **142**: 56–64

Suorsa M, Grieco M, Järvi S, Gollan PJ, Kangasjärvi S, Tikkanen M, Aro E-M (2013) PGR5 ensures photosynthetic control to safeguard photosystem I under fluctuating light conditions. *Plant Signal Behav* **8**: e22741

Suorsa M, Järvi S, Grieco M, Nurmi M, Pietrzykowska M, Rantala M, Kangasjärvi S, Paakkarinen V, Tikkanen M, Jansson S, et al (2012) PROTON GRADIENT REGULATION5 is essential for proper acclimation of *Arabidopsis* photosystem I to naturally and artificially fluctuating light conditions. *Plant Cell* **24**: 2934–2948

Tagawa K, Tsujimoto HY, Arnon DI (1963) Role of chloroplast ferredoxin in the energy conversion process of photosynthesis. *Proc Natl Acad Sci U S A* **49**: 567–572

Takahashi S, Badger MR (2011) Photoprotection in plants: a new light on photosystem II damage. *Trends Plant Sci* **16**: 53–60

Takahashi S, Milward SE, Fan D-Y, Chow WS, Badger MR (2009) How does cyclic electron flow alleviate photoinhibition in *Arabidopsis*. *Plant Physiol* **149**: 1560–1567

Tao R, Zhao Y, Chu H, Wang A, Zhu J, Chen X, Zou Y, Shi M, Liu R, Su N, et al (2017) Genetically encoded fluorescent sensors reveal dynamic regulation of NADPH metabolism. *Nat Methods* **14**: 720–728

Terai Y, Ueno H, Ogawa T, Sawa Y, Miyagi A, Kawai-Yamada M, Ishikawa T, Martua T (2020) Dehydroascorbate reductases and glutathione set a threshold for high-light-induced ascorbate accumulation. *Plant Physiol* **183**: 112–122

Terashima I, Funayama S, Sonoike K (1994) The site of photoinhibition in leaves of *Cucumis sativus* L. at low temperatures is photosystem I, not photosystem II. *Planta* **193**: 300–306

Tikhonov AN (2015) Induction events and short-term regulation of electron transport in chloroplasts: an overview. *Photosynth Res* **125**: 65–94

Tiwari A, Mamedov F, Grieco M, Suorsa M, Jajoo A, Styring S, Tikkanen M, Aro E-M (2016) Photodamage of iron–sulphur clusters in photosystem I induces non-photochemical energy dissipation. *Nat Plants* **2**: 16035

Trnka D, Engelke AD, Gellert M, Moseler A, Hossain MF, Lindenberg TT, Pedroletti L, Odermatt B, de Souza JV, Bronowska AK et al (2020) Molecular basis for the distinct functions of redox-active and FeS-transferring glutaredoxins. *Nat Commun* **11**: 3445

Vaseghi M-J, Chibani K, Telman W, Liebthal MF, Gerken M, Schnitzer H, Mueller SM, Dietz K-J (2018) The chloroplast 2-cysteine peroxiredoxin functions as thioredoxin oxidase in redox regulation of chloroplast metabolism. *eLife* **7**: e38194

Volpert A, Graff van Creveld S, Rosenwasser S, Vardi A (2018) Diurnal fluctuations in chloroplast GSH redox state regulate susceptibility to oxidative stress and cell fate in a bloom-forming diatom. *J Phycol* **54**: 329–341

Weise SE, Weber APM, Sharkey TD (2004) Maltose is the major form of carbon exported from the chloroplast at night. *Planta* **218**: 474–482

Wirtz W, Stitt M, Heldt HW (1982) Light activation of calvin cycle enzymes as measured in pea leaves. *FEBS Lett* **142**: 223–226

Yamamoto H, Shikanai T (2019) PGR5-dependent cyclic electron flow protects photosystem I under fluctuating light at donor and acceptor sides. *Plant Physiol* **179**: 588–600

Yamori W, Makino A, Shikanai T (2016) A physiological role of cyclic electron transport around photosystem I in sustaining photosynthesis under fluctuating light in rice. *Sci Rep* **6**: 1–12

Yoshida K, Hara A, Sugiura K, Fukaya Y, Hisabori T (2018) Thioredoxin-like2/2-Cys peroxidoxin redox cascade supports oxidative thiol modulation in chloroplasts. *Proc Natl Acad Sci U S A* **115**: E8296–E8304

Zhang Z, Jia Y, Gao H, Zhang L, Li H, Meng Q (2011) Characterization of PSI recovery after chilling-induced photoinhibition in cucumber (*Cucumis sativus* L.) leaves. *Planta* **234**: 883–889




Figure 1: Light-dependent redox modification of chloroplast-targeted roGFP2. (A) Confocal images of chl-roGFP2 Arabidopsis plants. Bars = 20 μm (B) Ratiometric analysis of chl-roGFP2 during rest, under oxidized (500 mM H_2O_2) and reduced (100 mM DTT) conditions using whole-plant imaging. The plants were excited with 405 nm, and 488 nm LEDs and fluorescence emission was collected using a 510 nm emission filter. For chlorophyll detection, a 405 nm LED light source and a 670 nm emission filter were used. Color-coded keys represent fluorescence intensities (below image) or 405/465 ratio values (to right of image). Bars = 2 cm. (C) The effect of light intensities on chl-roGFP2 reduction after application of H_2O_2 as examined by whole-plant imaging. Values represent means of 6 plants \pm SE. (D) and (E) The effect of DCMU on chl-roGFP2 oxidation using whole-plant imaging and plate-reader approaches. Oxidation of chl-roGFP2 was detected using whole-plant imaging. Bar = 2 cm (D) or a plate reader (E). Values for the degree of chl-roGFP oxidation (OxD) are presented as means of 7-8 plants \pm SE. Asterisks (*) mark significant differences between the ratio of the control/DCMU treatment (t-test, $P < 0.05$, Supplemental File 1). The roGFP data were acquired using the chl-TK_{TP}-roGFP2 line.




Figure 2: Diurnal changes in the degree of chl-roGFP2 oxidation under normal and high-light growth conditions. chl-roGFP2 fluorescence was monitored over a 24-hour period and chl-roGFP2 OxD values were calculated. Oxidation values were normalized to steady state values (as observed during the day at a constant light intensity of 120 $\mu\text{mol m}^{-2} \text{s}^{-1}$). Horizontal bars above graphs show light (yellow) and dark (black) periods. (A) Changes in chl-roGFP2 OxD over time from the light onset under normal light conditions. Values are presented as means of 8 plants \pm SE. (B) Changes in chl-roGFP2 OxD and normalized R_{400/485} of TKTP-iNAP4 over time during the nighttime. Results of TKTP-iNAP4 were normalized using the pH control sensor (TKTP-iNAPc) according to Lim et al. (2020) and values are presented as means of 12 plants \pm SE. (C) Oxidation and rereduction of chl-roGFP2 during dark-to-light transitions. Values are shown as means of 16-24 plants \pm SE. (D) Oxidation and rereduction of chl-roGFP2 during light-to-dark transitions, shown as means of 16-45 plants \pm SE. (E) Temporal changes in expression of chloroplast-targeted genes encoding antioxidant proteins and established ROS markers. Gene expression levels in plants exposed to HL (850 or 1700 $\mu\text{mol m}^{-2} \text{s}^{-1}$) were normalized to the gene expression in plants grown under normal light conditions (120 $\mu\text{mol m}^{-2} \text{s}^{-1}$) at each indicated time point after start of the HL illumination. All data are displayed as the log₂ ratio. The red and blue colors represent high or low expression levels, respectively. (F) Diurnal changes in chl-roGFP2 OxD in wild type (WT) plants exposed to HL. For each treatment, between 21 and 24 plants, divided into 3 independent plates, were tested. For clear presentation of the redox changes of roGFP2 over time, a “sliding window” approach for the data along the x-axis was taken, in which each data point represents the average of three plates (n=3). roGFP2 data were acquired using the chl-TKTP-roGFP2 line. Inset: Experimental design for the HL experiments. For a depiction of absolute roGFP2 OxD values and illustration of SE values see Supp.Fig.15 and Supp.Fig.16. (G) Diurnal changes in chl-roGFP2 Φ_{PSII} values in plants exposed to the same conditions as in (D). Φ_{PSII} values were derived from chlorophyll fluorescence analysis and represent means of 12 plants \pm SE.




Figure 3: Diurnal changes in chl-roGFP2 OxD under fluctuating light conditions. (A, D, G) Light irradiance during time over a 60-min period, extracted from the fluctuating light phase, is presented for the applied intervals of 1, 5 and 10 min. (B, E, H) Changes in chl-roGFP2 OxD values (ΔOxD) over time in *Arabidopsis* plants expressing chl-TKTP-roGFP2. After the start of the illumination period (yellow bar, $120 \mu\text{mol m}^{-2} \text{s}^{-1}$ for 3 h), the illumination regime was switched to fluctuating light alternating between HL ($1700 \mu\text{mol m}^{-2} \text{s}^{-1}$) and normal light ($120 \mu\text{mol m}^{-2} \text{s}^{-1}$) with different intervals. After 6 h fluctuating light (red bar), illumination continued continuously at $120 \mu\text{mol m}^{-2} \text{s}^{-1}$ for 7 h before the diurnal cycle ended with 8 h darkness (black bar). Oxidation values were normalized to steady state values (as observed during the day at constant light conditions of $120 \mu\text{mol m}^{-2} \text{s}^{-1}$). Each black line represents the oxidation dynamics recorded for an independent well plate which consists of 8 plants. Values represent means of these 8 plants \pm SE. (C, F, I) ΔOxD as a function of relative time (min) within each fluctuating light cycle. Values measured during the HL ($1700 \mu\text{mol m}^{-2} \text{s}^{-1}$) phases within each light cycle are presented in red. Values measured during the phases with normal light intensities ($120 \mu\text{mol m}^{-2} \text{s}^{-1}$) during each light cycle are presented in yellow. The presented box spans the interquartile range, and whiskers represent the minimum and maximum values (not including outliers). The median and mean of at least six plants are marked by a vertical line and x sign inside the box, respectively.




Figure 4: Diurnal changes in chl-roGFP2 OxD and Φ PSII under high-light and fluctuating light conditions in mutants with defects in photoprotective mechanisms. (A, B, C) Daily changes in chl-roGFP2 OxD under different light regimes in WT (A), *npq1* (B) and *pgr5* (C) plants expressing chl-PRX_{ATP}-roGFP2. The respective absolute ratio (400/485) data are provided in Supp. Fig.17. Oxidation values were normalized to steady state oxidation levels under normal growth light. The illumination regime was: 3 h normal light (yellow bar, 120 $\mu\text{mol m}^{-2} \text{s}^{-1}$), followed by 6 h HL (red bar, intensities as indicated by line color), 7 h normal light (120 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and 8 h darkness. Each treatment involved between 24 and 32 plants divided into four independent plates that were consolidated in a “sliding window” (n=4) display. Values represent means \pm SE. (D, E, F) Φ PSII for WT (D), *npq1* (E) and *pgr5* (F) plants exposed to the same light regimes as in A, B and C. Φ PSII values were derived from chlorophyll fluorescence measurements and represent means of 12 plants \pm SE. (G-I) Changes in chl-roGFP2 OxD during FL cycles with 10-min HL (1700 $\mu\text{mol m}^{-2} \text{s}^{-1}$) and 10 min normal light (120 $\mu\text{mol m}^{-2} \text{s}^{-1}$) for WT (G), *npq1* (H) and *pgr5* (I) plants. Values measured during the HL (1700 $\mu\text{mol m}^{-2} \text{s}^{-1}$) phases within each light cycle are presented in red. Values measured during the phases with normal light intensities (120 $\mu\text{mol m}^{-2} \text{s}^{-1}$) during each light cycle are presented in yellow. The presented box spans the interquartile range, and whiskers represent the minimum and maximum values (not including outliers). The median and mean of at least six plants are marked by a vertical line and x sign inside the box, respectively.




Figure 5: Diurnal changes in chl-roGFP2 OxD and Φ_{PSII} during one day of high-light regime followed by two days of normal light conditions in *pgr5* and *gl1* plants. (A, C) Changes in chl-roGFP2 OxD over time from the experiment onset. chl-roGFP2 fluorescence was monitored throughout the experiment and chl-roGFP2 OxD values were calculated. Oxidation values were normalized to steady state oxidation levels. Each treatment involved 32 plants divided into four independent plates that were consolidated in a “sliding window” ($n=4$) display. Values represent means \pm SE for *pgr5* (A) and *gl1* (C). roGFP data were acquired using the chl-PRX α_{ATP} -roGFP2. (B) Φ_{PSII} over time from the experiment onset. Φ_{PSII} values were derived from chlorophyll fluorescence measurements and represent means of 12 plants \pm SE for *pgr5* (B) and *gl1* (D). The color bar denotes the light conditions: black – night, yellow – normal light, red – high light